

A 3D Automated Classification Scheme for the TAUVE X data pipeline

Archana Bora^{1,2*}, Ranjan Gupta¹, Harinder P. Singh³, Jayant Murthy⁴
Rekshesh Mohan⁴ and Kalpana Duorah²

¹*IUCAA, Post Bag 4, Ganeshkhind, Pune-411007, India*

²*Department of Physics, Gauhati University, Guwahati 781014, India*

³*Department of Physics & Astrophysics, University of Delhi, Delhi 110007, India*

⁴*Indian Institute of Astrophysics, Koramangla, Bangalore 560034, India*

Received on 15/6/2007; Accepted on 23/11/2007

ABSTRACT

In order to develop a pipeline for automated classification of stars to be observed by the TAUVE X ultraviolet space Telescope, we employ an artificial neural network (ANN) technique for classifying stars by using synthetic spectra in the UV region from 1250Å to 3220Å as the training set and International Ultraviolet Explorer (IUE) low resolution spectra as the test set. Both the data sets have been pre-processed to mimic the observations of the TAUVE X ultraviolet imager. We have successfully classified 229 stars from the IUE low resolution catalog to within 3-4 spectral sub-class using two different simulated training spectra, the TAUVE X spectra of 286 spectral types and UVBLUE spectra of 277 spectral types. Further, we have also been able to obtain the colour excess (i.e. $E(B-V)$ in magnitude units) or the interstellar reddening for those IUE spectra which have known reddening to an accuracy of better than 0.1 magnitudes. It has been shown that even with the limitation of data from just photometric bands, ANNs have not only classified the stars, but also provided satisfactory estimates for interstellar extinction. The ANN based classification scheme has been successfully tested on the simulated TAUVE X data pipeline. It is expected that the same technique can be employed for data validation in the ultraviolet from the virtual observatories. Finally, the interstellar extinction estimated by applying the ANNs on the TAUVE X data base would provide an extensive extinction map for our galaxy and which could in turn be modeled for the dust distribution in the galaxy.

Key words: ISM: dust – extinction methods: data analysis – space vehicles: instruments – astronomical databases: miscellaneous – ultraviolet: general

1 INTRODUCTION

Tel-Aviv University Ultra-Violet Experiment (TAUVE X) is an Indo-Israeli Ultraviolet Imaging space mission that will image large parts of the sky in the wavelength region between 1300 and 3200Å. The instrument consists of three equivalent 20-cm UV imaging telescopes with a choice of filters for each telescope. Each telescope has a field of view of about 54' and a spatial resolution of about 6" to 10", depending on the wavelength. TAUVE X will be launched into a geostationary orbit as part of Indian Space Research Organization's GSAT-4 mission in April 2008.

Observations will be available using filters in five UV bands:

- (i) BBF : Broadband filter (1300–3300Å)
- (ii) SF1 : Intermediate band filter 1 (1250–2250Å)
- (iii) SF2 : Intermediate band filter 2 (1800–2600Å)
- (iv) SF3 : Intermediate band filter 3 (2100–3100Å)
- (v) NBF : Narrowband filter (2000–2400Å)

Figure 1 shows the response curves for each of the TAUVE X filters in units of Effective Area cm^2 .

The TAUVE X mission will have added advantages as compared to other earlier UV missions like the TD satellites and GALEX etc. The estimation of the slope R_v of the interstellar extinction curve with a greater sensitivity, will allow to construct deeper maps of the UV sky. Further, TAUVE X and TD satellites would complement each other by having a

* E-mail: archana@iucaa.ernet.in; rag@iucaa.ernet.in; hpsingh@physics.du.ac.in; jmurthy@yahoo.com reks@iia.res.in and kalpanaduorah@yahoo.com




Figure 1. Filter response of the five filters of TAUVE X

total of six data points for the interstellar extinction curve for their common sources (see Maheshwar et al. 2007).

TAUVE X will mostly operate in scanning mode, since it will be mounted on GSAT-4, a geosynchronous satellite. The FOV will be scanning a strip of the sky with constant declination and a limiting magnitude of 19 (Murthy, 2003). A few years of successful run of the mission will record more than a million UV point sources apart from galaxies, QSO's and the UV background. The need for an automated classification pipeline for the stellar sources that is repeatable and fast is, therefore, immense.

The Artificial Neural Network (ANN) based schemes are now being routinely used to classify spectra from large spectral data bases (Gulati et al. 1994, Singh et al. 1998, 2006, Valdes et al. 2004, Bailer-Jones 2002, Gupta et al. 2004) for the purpose of sorting these large spectral data base into groups of main spectral types (O, B, A, F, G, K and M) and sub-classes. Further, these schemes can also be used for obtaining stellar fundamental atmospheric parameters (Gulati et al. 1997 a,b). Of these Gulati et al. (1997b) is of particular interest, since it was shown that ANNs can determine the colour excess, i.e. $E(B-V)$ in units of magnitudes, as an additional parameter when applied to the IUE spectral data base.

The current work has used the ANN based tools for classifying the IUE spectral data base (reduced to the TAUVE X band data) in terms of the spectral types and also hierarchically estimated the color excess using this tool. It is worth noting that whereas the Gulati et al. (1997b) used the IUE full spectra for spectral type classification and estimation of colour excess, the present work uses the simulated band data as expected from the TAUVE X satellite and even with this limitation, the neural network scheme has been able to assign the spectral classes and also obtain reddening estimate to satisfactory levels.

In the next section, we describe the generation and pre-

processing of the simulated TAUVE X data that is used for training of the neural network as well as the processing of the IUE spectral data which is used as the test set. In Section 3, we describe results of the ANN classification scheme as well as the color excess determination. In Section 4, we present important conclusions of the study.

2 ANN ARCHITECTURE, GENERATION OF SIMULATED DATA AND ANN TRAIN AND TEST SETS

Following sub-sections describe the ANN architecture, simulated data generation and the ANN train and test sets.

2.1 ANN architecture

The ANN architecture considered here is an supervised one with a minimum configuration of three layers, i.e., (1) Input layer where the patterns are read (2) Hidden layer where the information is processed from the input layer (3) Output layer where the output patterns are rendered (see Bailer-Jones et al. 2002 for a review). The hidden layer can have several nodes which inter-connect the input and output layers with each connection with its designated connection weight. We have used a back-propagation algorithm (Gulati et al. 1994, 1997a,b, Singh et al. 1998) with 2 hidden layers of 64 nodes each and this scheme requires a training session where the ANN output and the desired output get compared after each iteration and the connection weights get updated till the desired minimum error threshold is reached. At this stage the network training is complete and the connection weights are considered frozen. The next stage is the testing session where the test patterns are fed to the network and output is the classified spectral pattern or color excess in terms of the training sets.

In the actual post launch of TAUVE X when the real data will be available, the scheme applied to estimate the colour excess will have to run the ANN in two stages i.e. in a hierarchal manner such that, the first stage classifies the test set (IUE data base or the expected TAUVE X data base) into the spectral classes and then a second ANN stage performs the colour excess estimation.

2.2 Simulated data generation

We have used two independent sources to generate the training sets of spectra with solar type stars with $[M/H] = 0$. One is the stellar flux calculator from TAUVE X web-site (<http://tauvex.iap.res.in/htmls/tools/fluxcalc/>) containing 286 spectro-luminosity classes and the other is the UVBLUE fluxes (Rodriguez-Merino et al. 2005) (<http://www.bo.astro.it/~eps/uvblue/uvblue.html>). Based on the spectral type and luminosity class of a star, the TAUVE X calculator derives the effective temperature and surface gravity using the calibration of Allen (2000), Colina (1995) and Lang (1982) and calculates the spectral energy distribution for each star using appropriate Kurucz model available on the webpage <http://kurucz.harvard.edu/> (see Sujatha et al. 2004). We have used the information from Allen (2000), Erika Böhm-Vitense (1981), Johnson (1966), Ridgway et al. (1980), Alonso et al. (1999) and Bertone et

al. (2004) for matching the parameter space of UVBLUE to spectral-types and luminosity classes. Both the sources provide sets of theoretical fluxes (based on Kurucz model atmospheres) in the UV region. These fluxes need to be processed via a common flux integration programme provided at the TAUVE X tools site to form two sets of band data (each having four fluxes corresponding to the four TAUVE X bands) and they constitute the simulated band data set for the ANN training sets.

We have also obtained two sets of fluxes (with 50\AA resolution and 40 data bins covering the spectral region of $1250\text{--}3220\text{\AA}$) aimed at preparing the ANN tools for another Indian scientific mission satellite ASTROSAT (<http://www.rri.res.in/astrosat/>) which will have gratings to provide slit-less spectra for spatially resolved stars. It will also prepare us for the future GAIA mission (<http://gaia.esa.int/science-e/www/area/index.cfm?fareaid=26>).

2.3 ANN train and test sets

While making the train and test sets, one has to ensure that the number of spectral fluxes at the respective wavelengths and the starting/ending wavelengths are identical. Also the spectral resolution needs to be same and for this, the spectral libraries had to be convolved with appropriate gaussian functions to bring them at par with each other. The fluxes are normalized to unity with respect to maximum flux in each spectrum before sending to the ANN inputs. The spectra for 286 TAUVE X spectral types generated in the range $1250\text{--}3200\text{\AA}$ have a resolution of 10\AA which we have degraded to 50\AA . The resolution of 277 UVBLUE spectral types have been degraded similarly (using the relevant codes provided on the UVBLUE library web site). These sets of data are then reddened (using the observed extinction curve of Seaton, 1979) in the range of 0.00-1.00 mag, for preparing the training sets for the two stages of the hierarchal scheme viz. the separation of the different spectral types and the evaluation of reddening values. Below we provide the details of the procedure adopted for generating the training sets for the two stages:

• Generating data set for Spectral Type determination:

In the first stage, reddening values are added in step sizes of 0.20 magnitudes to the simulated data. The 0.20 step is chosen for the computational convenience. For example, the TAUVE X data consists of 286 different classes with 58 spectral types, each having 5 luminosity classes (except for O6.5V). If one wants to classify the Spectral type, Luminosity class and the reddening value in a single run; reddening these 286 data sets with reddening value from 0.00-1.00, even at a step of 0.1 leads to $286 \times 11 = 3146$ number of distinct classes. However, this is not possible with our current computational facilities and the present version of our ANN. Instead, we go for the hierarchal scheme by first merging all the luminosity classes. For example, instead of considering O3I-O3V as five separate classes, the ANN will be trained to learn all the five different patterns as single O3 spectral type only, though the variation in all the five spectra still go as input to the ANN. The process thus reduces the number of distinct classes from 286 to only 58 classes, making the computation fast. When the learning process is completed,

ANN can separate different Spectral types, thus making it possible to find out the reddening values in the next stage.

• Generating data set for Reddening evaluation:

In the second stage, reddening values are added in step sizes of 0.05 to the simulated data. The separation of the available spectra into different groups O, B, A, F, G, K etc. in the first stage, makes it possible to select this finer step size of 0.05. In our work we have not classified the luminosity classes separately, however, this can be done easily by adding one more stage in the hierarchal scheme.

A sample of normalized simulated spectra of different spectral types is shown in Fig. 2. Their integrated fluxes in the four TAUVE X bands, NBF, SF1, SF2 and SF3, have been computed using respective filter response curves of Fig.1. Fig. 3 shows the residuals obtained by subtracting IUE fluxes from the corresponding TAUVE X simulated fluxes. The discrepancies observed in these figures could be due the following reasons:

In the early-type stars i.e. O and B, the main discrepancy between observed and theoretical is near 1500\AA . This is a consequence of the physical origin of the CIV line, which gets strongly affected by stellar winds and mass-loss processes in massive stars. For F-type stars the metallic features at 2400\AA (Fe III), 2500\AA (Fe I/Si I), 2800\AA (Mg II) are more enhanced in the simulated spectra. For G-type stars the chromospheric activities increases and thus triggers prominent Mg core emissions which are not seen in the simulated spectra. The chromospheric activities are not accounted for in the Kurucz's model (Rodriguez-Merino et al. 2005). The discrepancy is more clearly visible in the band integration of the fluxes of late type stars in Fig. 3.


The final training set thus contains (a) the spectra in the form shown in Fig. 2 and (b) 4 flux values in the 4 bands of TAUVE X in the form shown in Fig. 3 – for each of the 286 TAUVE X spectra (277 spectra for the UVBLUE case) with reddening in the range of 0.00 to 1.00 mag with a step of 0.2 mag. Fig. 4 shows a block diagram of the flow chart for preparing these two training sets for spectral type classification.

The test spectra were taken from the IUE low resolution spectra: reference atlas, normal stars, ESA SP-1052 by Heck et al. (1984) which contains 229 low-dispersion flux calibrated spectra of O to K spectral type obtained by the IUE satellite. The spectra were trimmed to $1250\text{--}3220\text{\AA}$. The original resolution of 6\AA of IUE spectra was convolved by a Gaussian function to produce a degraded resolution of 50\AA . Fig. 5 shows the block diagram of the flow chart for generating this IUE test set for spectral classification. The Fig. 6 shows a block diagram for the flow chart for creating the train set for extinction classification and Fig. 7 shows the corresponding block diagram of the flow chart for creating the IUE test set.

Table 1 shows the number of spectra per spectral type used in this analysis. The numbers in the 2nd and 3rd column are the basic sets for training sessions of the ANN. The hierarchal ANN scheme used by us works in two stages viz. Ist stage performs the spectral type classification and for this these numbers get multiplied by 6 and in the IInd stage which performs the color excess classification, they get multiplied by 21. Further, in order to have an uniform number

Table 1. Number of Spectra for each data set according to the spectral types.

Spectral Class	TAUVEX	UVBLUE	IUE
O	36	36	42
B	50	41	115
A	50	50	48
F	50	50	20
G	50	50	3
K	50	50	1

**Figure 2.** IUE and TAUVEK Simulated fluxes for 6 sample stars at a resolution of 50 Å.

of spectra per spectral type, classes which have just one example are duplicated during the training session.

3 RESULTS OF THE ANN CLASSIFICATION

The results of spectral classification are depicted in the Fig. 8. The numbers on the axes of this figure refer to the spectral coding which is briefly described as follows:


Main Spectral Type: O = 1000, B = 2000, A = 3000, K = 6000,

Sub-Spectral Type: O1 = 1100, O2 = 1200, O9 = 1900,

Luminosity Class: I = 1.5, II = 3.5, III = 5.5, IV = 7.5 and V = 9.5.

For example, Sun is a G2V star and hence its code will be 5209.5. A Classification error of 500 implies that a G2 star can, at worse, be classified either as F7 or G7 spectral type.

Figure 9 shows the scatter plots for pre-classified IUE stars (in O, B, A and F spectral types) for UVBLUE fluxes

**Figure 3.** Integrated IUE and Simulated TAUVEK fluxes for the same 6 sample stars in NBF, SF1, SF2 and SF3 filters along with the residues in the corresponding lower panels.

with their colour excess estimates σ in units of magnitudes. Figure 10 shows the scatter plots for pre-classified IUE stars (in O, B, A and F spectral types) for UVBLUE bands with their colour excess estimates σ in units of magnitudes. Figures 11 & 12 show the corresponding classification results for TAUVEK fluxes and bands respectively. In these 3D scatter plots, the 'Cat' and 'ANN' denote the catalog and ANN classes respectively. Further, the vertical axis in the plots gives the number of stars (N) present for a particular color excess value and are re-scaled as the square root of the actual number (i.e. $N^{1/2}$) for better representation; otherwise in the cases where this number is large, the corresponding points for single stars would look too small on the plots.

It is important to see that in the spectral classification scheme, the outliers in the all the four panels of Fig. 8 belong to G and K type, they being misclassified as the F type stars. This can be attributed to the discrepancies mentioned in section 2.3. In the two exceptional cases G8 gets classified as O2 type in FLUX UVBLUE panel whereas A2 gets classified as K3 in FLUX TAUVEK panel. The misclassification of G8 as O2 may be because as G8 IUE spectra shows a moderate UV excess compared to the theoretical one as mentioned in Rodriguez-Merino et al. (2005).




Figure 4. A block diagram showing the flow chart for creating the ANN train set for spectral classification with TAUVE X and UVBLUE simulated sources.

From the Figs. 9, 10, 11 & 12 we see an overall colour excess estimate accuracy in the range of 0.20 in the worst case of F-Type spectra with bands to 0.06 in the best case for B-Type spectra with bands. The results with bands show better accuracies in comparison to the fluxes which may indicate that band data is a better estimator for colour excess than the fluxes.

The ANN inputs take most of the information in terms of absorption features which are embedded in the full range of spectral fluxes (or the integrated fluxes in the band data) for performing the classification. This information is available for the hot stars like O, B and A but lacks in F or later spectral types. Due to this reason, the ANNs do not provide a good estimate of reddening for these late type stars. Thus we have not estimated the colour excess for the G and K Type IUE spectra (the 3 nos of G Type and 1 no of K Type of the IUE test set mentioned in Table 1 have no reddening). Table 2 summarizes the results for both spectral type classification and colour excess estimation.

4 CONCLUSIONS

Till now several studies have demonstrated that the artificial neural network schemes can reliably and successfully classify stellar spectral data as well as extract fundamental stellar parameters in the visible region. The extension of applicability of this scheme to UV region has been less prevalent mainly because of non-availability of abundant data in this




Figure 5. A block diagram showing the flow chart for creating the ANN test set for corresponding to the Figure 4.

region. Nevertheless, some attempts have been made in the past to automate the process of classification of spectral data from the IUE satellite. In this paper, we have demonstrated that the artificial neural networks can be successfully employed to classify stellar photometric (band) data.

We have shown that the ANN tools developed by us can successfully classify the 229 IUE spectra reduced to the four TAUVE X bands to an accuracy in the range of 3-4 sub-spectral types. We have also estimated the colour excess for the hot stars (O, B and A types) to an accuracy of up to 0.1 magnitudes in terms of $E(B-V)$ colours. Thus, even with the limitation of data from just photometric bands, ANNs have not only classified the stars, but also provided satisfactory estimates for interstellar extinction.



We hope that our automated pipeline will be used extensively to extract and validate data from virtual observatories as well as for the upcoming satellite data base expected from the TAUVE X and also the ASTROSAT and GAIA missions where one will be able to provide the interstellar extinction maps of our galaxy and which in turn could be modeled for dust distribution (Vaidya et al. 2001, Gupta et al. 2005, Vaidya et al. 2007).

ACKNOWLEDGMENTS

This work is supported by a Grant from Indian Space Research Organization under its RESPOND scheme. We would like to acknowledge an email correspondence with Prof. E.

Table 2. Summary of Classification results.

Spectral Classification Error σ (sub-spectral type)				
Simulated Source:	TAUVEX		UVBLUE	
	Flux	Band	Flux	Band
	3.97	3.84	3.77	3.39
Colour Excess E(B-V) Error σ (magnitudes)				
O-Type	0.10	0.09	0.11	0.09
B-Type	0.09	0.07	0.08	0.06
A-Type	0.10	0.08	0.14	0.09
F-Type	0.10	0.16	0.18	0.20

**Figure 6.** A block diagram showing the flow chart for creating the ANN train set for extinction classification for both simulated sources i.e. TAUVEK and UVBLUE.**Figure 7.** A block diagram showing the flow chart for creating the ANN test set for extinction classification corresponding to the Fig. 6.

Bertone, for providing the relevant references which helped us in matching the UVBLUE parameter space to spectral types and luminosity classes. AB thanks Dr. Kalpana Duorah for inspiring her to undertake this project.

REFERENCES

- Allen 2000, *Astrophysical Quantities*, 4th ed.
 Alonso, A., Arribas, S. and Martinez-Roger, C., 1999, *A&AS*, 140, 261

- Bailer-Jones, C. A. L. 2002, in *Automated Data Analysis in Astronomy*, ed. R. Gupta, H. P. Singh, & C. A. L. Bailer-Jones (New Delhi: Narosa), 83
 Bailer-Jones, C. A. L., Gupta, R., & Singh, H. P. 2002, in *Automated Data Analysis in Astronomy*, ed. R. Gupta, H. P. Singh, & C. A. L. Bailer-Jones (New Delhi: Narosa), 51
 Bertone, E., Buzzoni, A., Rodriguez-Merino, L. H. and Chavez, M., 2004, *AJ*, 128, 829
 Colina, L., 1995, *CDBS Kurucz Stellar Atmosphere, Instrument Science Report SCS/CAL-006 (STScI/OSG)*
 Erika B  hm-Vitense, 1981, *Ann. Rev. Astron. Astrophys.*,




Figure 8. Scatter plots of classification of the 229 IUE stars with TAUVE X bands and fluxes and with UVBLUE bands and fluxes. The classification accuracy values σ are shown for each case in units of sub-spectral types.




Figure 9. Scatter plot of classification of 229 IUE stars (pre-classified into O, B, A and F spectral types) with TAUVE X fluxes and UVBLUE fluxes for colour excess estimates. The classification accuracy values σ are shown for each case in units of E(B-V) magnitudes.




Figure 10. Scatter plot of classification of 229 IUE stars (pre-classified into O, B, A and F spectral types) with UVBLUE bands for colour excess estimates. The classification accuracy values σ are shown for each case in units of E(B-V) magnitudes.




Figure 11. Scatter plot of classification of 229 IUE stars (pre-classified into O, B, A and F spectral types) with TAUVE X fluxes for colour excess estimates. The classification accuracy values σ are shown for each case in units of E(B-V) magnitudes.

295, 318

Gulati, R. K., Gupta, R., Gothoskar, P., & Khobragade, S. 1994, ApJ, 426, 340

Gulati, R. K., Gupta, R., & Rao, N. K. 1997a, A&A, 322, 933

Gulati, R. K., Gupta, R., & Singh, H. P. 1997b, PASP, 109, 843

Gupta, R., Singh, H. P., Volk, K. and Kwok, S., 2004, ApJS, 152n2, 201

Gupta, R., Mukai, T., Vaidya, D. B., Sen, A. K. and Okada, Y. 2005, A&A, 441, 555

Heck, A., Egret, D., Jaschek, M., Jaschek, C. 1984, IUE Low-Resolution Spectra: a Reference Atlas – Part I, Normal Stars, ESA SP-1052

Johnson, H. L., 1966, Ann. Rev. Astron. Astrophys., 4, 193




Figure 12. Scatter plot of classification of 229 IUE stars (pre-classified into O, B, A and F spectral types) with TAUVEK fluxes for colour excess estimates. The classification accuracy values σ are shown for each case in units of $E(B-V)$ magnitudes.

- Lang, K. R., 1982, *Astrophysical Data: Planets and Stars*, Springer-Verlag
- Maheshwar, G., Muthu, C., Sujatha, N. V., Pandey, G., Bhatt, H. C., Kameswara Rao, N. and Murthy, J., 2007, *BASI*, 35, 233
- Murthy, J. 2003, *BASI*, 31, 243
- Ridgway, S. T., Joyce, R. R., White, N. M. and Wing, R. F., 1980, *ApJ*, 235, 126
- Rodriguez-Merino, L. H., Chavez, M., Bertone, E., Buzoni, A. 2005, *ApJ*, 626, 411 (UVBLUE)
- Seaton, M. J., 1979, *MNRAS*, 187, 73
- Singh, H. P., Gulati, R. K., Gupta, R. 1998, *MNRAS*, 295, 312
- Singh, H. P., Yuasa, M., Yamamoto, N., Gupta, R. 2006, *PASJ*, 58, 177
- Sujatha, N. V., Chakraborty, P., Murthy, J. and Henry, R. C., 2004, *BASI*, 32, 151
- Vaidya, D.B., Gupta, R., Dobbie, J. S., Chylek, P. 2001, *A & A*, vol. 375, 584
- Vaidya, D.B., Gupta R., Snow, T. P. 2007, *MNRAS*, 371, 791
- Valdes, F., Gupta, R., Rose, J. A., Singh, H. P., Bell, D. J. 2004, *ApJS*, 152, 251

